

Exact solutions for a Dirac electron in an exponentially decaying magnetic field

Tarun Kanti Ghosh

Department of Physics, Indian Institute of Technology-Kanpur, Kanpur 208 016, India

(Dated: November 30, 2009)

We consider a Dirac electron in the presence of an exponentially decaying magnetic field. We obtain exact energy eigenvalues with a zero-energy state and the corresponding eigenfunctions. We also calculate the probability density and current distributions.

PACS numbers: 03.65.Pm, 73.20.-r, 81.05.Uw, 71.70.Di

The experimental realization of graphene, two-dimensional (2D) sheet of graphite [1, 2], and of the massless Dirac nature of its electron low-energy spectrum [3, 4] has given rise to a tremendous interest in this field (for recent reviews, see [5, 6]). The energy spectrum which goes linearly with the momentum and the specific density of states of the Dirac electrons [7] enabled the study of experimentally chiral tunneling and the Klein paradox in graphene [8]. This also leads to the anomalous Landau level spectrum in a uniform magnetic field, which gives rise to the half-integer quantum Hall effect [3, 9].

The discovery of the half-integer quantum Hall effect and the zero-energy Landau level [9] stimulated lot of theoretical research interest on Dirac electrons in uniform as well as non-uniform magnetic fields. For example, the Dirac-Weyl equation has been solved numerically for a single electron in a step-like magnetic field, magnetic barrier and polytropic magnetic field with $B = B_0 y^\gamma$, where $\gamma > 0$ [10, 11, 12, 13]. We should also mention that the Schrödinger equation has been solved numerically for spinless 2D electrons in a linearly varying magnetic field [14], step-like magnetic field, magnetic barrier [15, 16, 17]. An analytical solution has been given for an electron in the presence of an exponentially decaying magnetic field [18].

In this work, we solve analytically for the Dirac electron in graphene in the presence of an exponentially decaying magnetic field $B = B_0 e^{-\lambda x}$.

The Hamiltonian of the massless electrons in graphene near one of the Dirac K points is described by a two-component Dirac-Weyl equation

$$H = v_F \sigma \cdot \mathbf{p}, \quad (1)$$

where $v_F \approx 10^6$ m/sec is the Fermi velocity, $\sigma = \{\sigma_x, \sigma_y\}$ is the 2×2 Pauli matrices and $\mathbf{p} = -i\hbar\nabla$ is the two-dimensional momentum operator. The pseudo-relativistic dispersion relation with Fermi velocity arises is due to the sublattice structure: the basis of the graphene honeycomb lattice contains two carbon atoms, giving rise to an isospin degree of freedom. In the presence of an external magnetic field ($\mathbf{B} = \nabla \times \mathbf{A}$) perpendicular to the graphene plane, the Hamiltonian of the single Dirac electron ($-e$) is $H = v_F \sigma \cdot (\mathbf{p} + e\mathbf{A})$.

We consider an electron in the graphene sheet in the presence of a non-uniform magnetic field $\mathbf{B}(x, y) =$

$\{0, 0, B_0 e^{-\lambda x} \hat{z}\}$ perpendicular to the $x - y$ plane. In the Landau gauge, the corresponding vector potential is $\mathbf{A}(x, y) = \{0, -(1/\lambda)B_0 e^{-\lambda x} \hat{y}, 0\}$. When $\lambda \rightarrow 0$, the non-uniform magnetic field becomes a constant magnetic field. The time-independent Dirac-Weyl equation is

$$v_F \sigma \cdot (\mathbf{p} + e\mathbf{A})\Psi(x, y) = E\Psi(x, y). \quad (2)$$

Here, $\Psi(x, y) = \{\Psi_+(x, y), \Psi_-(x, y)\}^T$ is the two-component wavefunction and T denotes the transpose of the column vector. Due to the translation invariance in the y -direction, the longitudinal momentum k_y is a conserved quantity. One can parameterize the wavefunction as $\Psi(x, y) = e^{ik_y y} \{\psi_+(x), \psi_-(x)\}^T$. From equation (2), one can get two coupled equations for ψ_+ and ψ_- as given below:

$$-i\hbar v_F \left[\frac{\partial}{\partial x} + (k_y + \frac{e}{\hbar} A_y) \right] \psi_-(x) = E\psi_+(x) \quad (3)$$

and

$$-i\hbar v_F \left[\frac{\partial}{\partial x} - (k_y + \frac{e}{\hbar} A_y) \right] \psi_+(x) = E\psi_-(x). \quad (4)$$

Using two coupled equations. (3) and (4), we get Schrödinger-like decoupled equations for ψ_+ and ψ_- as

$$[-(\hbar v_F)^2 \frac{\partial^2}{\partial x^2} + V_+(x)]\psi_+ = E^2\psi_+ \quad (5)$$

and

$$[-(\hbar v_F)^2 \frac{\partial^2}{\partial x^2} + V_-(x)]\psi_- = E^2\psi_-. \quad (6)$$

Here, the effective potentials $V_\pm$ are given by

$$V_\pm = (\hbar v_F)^2 \left[\pm \frac{e}{\hbar} \frac{\partial}{\partial x} A_y(x) + (k_y + \frac{e}{\hbar} A_y)^2 \right]. \quad (7)$$

Here, the first term on the right hand side of the above equation is the Zeeman-like term due to the isospin degree of freedom in presence of the magnetic field. Define the magnetic length scale as $l(x) = \sqrt{\hbar/eB(x)}$.

As seen in figure 1, the effective potentials $V_\pm(x)$ have the form of an asymmetric quantum wells formed by the exponentially decaying magnetic field. It is well known that such a well can have a bound state if the well is

sufficiently deep. Figure 1 will be discussed in more detail later on.

Following the references. [18, 19], we introduce two new dimensionless variables:

$$\xi(x) = \frac{1}{(l(x)\lambda)^2} = \frac{eB_0 e^{-\lambda x}}{\hbar\lambda^2} \quad (8)$$

and

$$\xi_0 = \frac{|k_y|}{\lambda} \equiv \frac{eB_0 e^{-\lambda x_0}}{\hbar\lambda^2} = \frac{1}{(l(x_0)\lambda)^2}. \quad (9)$$

Here, $l(x_0) = \sqrt{\hbar/eB(x_0)}$ is the magnetic length scale for a given value of $x = x_0$ which depends on the conserved wavevector k_y through equation (8). Also, ξ varies from 0 to ∞ when x varies from ∞ to $-\infty$.

In the new variables, equations. (5) and (6) reduce to

$$\left[\frac{d^2}{d\xi^2} + \frac{1}{\xi} \frac{d}{d\xi} - \frac{\beta^2}{\xi^2} + \frac{2\xi_0 \mp 1}{\xi} - 1 \right] \psi_{\pm} = 0, \quad (10)$$

where $\beta^2 = \xi_0^2 - \epsilon^2$ and $\epsilon = E/(\hbar v_F \lambda)$. The behavior for small and large ξ suggests that the general solution can be written as $\psi_{\pm}(\xi) \sim \xi^{\beta} e^{-\xi} w_{\pm}(\xi)$. Inserting this ansatz in the previous equation, we get for $w_{\pm}(z = 2\xi)$

$$\left[z \frac{d^2}{dz^2} + (\gamma - z) \frac{d}{dz} - \alpha_{\pm} \right] w_{\pm}(z) = 0 \quad (11)$$

which has the form of a confluent hypergeometric equation, where $\gamma = 2\beta + 1$ and $\alpha_{\pm} = -\xi_0 + \beta + \frac{1}{2} \pm \frac{1}{2}$. Two linearly independent solutions can be chosen as $F[\alpha_{\pm}, \gamma; z]$ and $U[\alpha_{\pm}, \gamma; z]$, so the general solution can be written as $w_{\pm}(z) = A_1 F[\alpha_{\pm}, \gamma; z] + A_2 U[\alpha_{\pm}, \gamma; z]$. Here, F and U are the first and second kind of confluent hypergeometric functions, respectively. However, U is not regular at the origin and has to be discarded. The requirement of normalizability implies that the solution is acceptable if $\alpha_{\pm}$ is negative integer $\alpha_{\pm} = -\nu, \nu = 0, 1, 2, \dots$. This constraint produces the quantization of the energy:

$$E_{\pm} = \frac{\hbar v_F \lambda}{2} \sqrt{(2\xi_0)^2 - (2\xi_0 - (2\nu + 1 \pm 1))^2}. \quad (12)$$

The energy eigenvalues are then conveniently written as

$$E_n = \hbar v_F \lambda \sqrt{(\xi_0)^2 - (\xi_0 - n)^2}. \quad (13)$$

For E_+ , $\nu = n - 1$ and for E_- , $\nu = n$. For $n = 0$ and E_+ , $\nu = -1$, but ν can not be negative. For $n = 0$ and E_- , $\nu = 0$. Therefore, the $n = 0$ state is not degenerate. For $n = 0$, the solution w_+ does not exist. We will consistently incorporate this fact by defining $w_{n-1} = 0$. The corresponding wave function is given by

$$w_{\pm}(2\xi) = {}_1F_1[-(n - \frac{1}{2} \mp \frac{1}{2}), 2\beta + 1, 2\xi], \quad (14)$$

where ${}_1F_1[-n, \alpha, x]$ is the confluent hypergeometric function.

Equation (13) can be re-written as

$$E_n^2 = \hbar^2 \omega_c^2(x_0) 2n(1 - \frac{n}{2\xi_0}), \quad (15)$$

where the local cyclotron frequency is $\omega_c(x_0) = (v_F)/l(x_0)$. Note that when $\bar{\lambda} \rightarrow 0$ ($\xi_0 \rightarrow \infty$), the energy eigenvalues reduce to the well known relativistic Landau level structure for uniform magnetic field: $E_n = (\hbar v_F/l(x_0))\sqrt{2n}$.

When $\bar{\lambda}$ is very large, the effect of inhomogeneous magnetic field vanishes and the Dirac electron feels no effective potential. It behaves like a free particle. In this limit, the energy spectrum becomes $E_n \propto n$ which agrees with the known result for the free Dirac electron [20].

The complete normalized wavefunctions can be written as

$$\Psi_+^{(n)} = \frac{e^{ik_y y}}{\sqrt{2L_y l(x_0)}} (2\xi_0)^{\beta} e^{-\bar{\lambda}\beta X} e^{-\xi_0 e^{-\bar{\lambda}X}} w_{n-1} \quad (16)$$

and

$$\Psi_-^{(n)} = \frac{ic_n e^{ik_y y}}{\sqrt{2L_y l(x_0)}} (2\xi_0)^{\beta} e^{-\bar{\lambda}\beta X} e^{-\xi_0 e^{-\bar{\lambda}X}} w_n, \quad (17)$$

where L_y is the length of the system along the y axis, $c_0 = \sqrt{2}$ and $c_{n>0} = 1$. Also, $X = (x - x_0)/l(x_0)$, $\bar{\lambda} = \lambda l(x_0)$, and w_n is given by

$$w_n = \sqrt{\frac{\Gamma[2\beta + n + 1]\bar{\lambda}}{\Gamma[2\beta + 1]\Gamma[2\beta]\Gamma[n + 1]}} {}_1F_1[-n, 2\beta + 1, 2\xi_0 e^{-\bar{\lambda}X}]. \quad (18)$$

We have also checked that the wavefunctions (16) and (17) reduce to that of the Dirac Landau level for constant magnetic field (*i.e.* a Hermite polynomial multiplied with a Gaussian factor) when $\bar{\lambda} \rightarrow 0$. The phase factor i in the lower component wavefunction (17) is obtained from equation (4), which is crucial for calculating the probability current density.

The effective potential (7) can be re-written as

$$V_{\pm} = (\hbar\omega_c(x_0))^2 [\pm e^{-\bar{\lambda}X} + \xi_0(1 - e^{-\bar{\lambda}X})^2]. \quad (19)$$

For weak-inhomogeneity ($\bar{\lambda} \ll 1$), the effective potentials are almost symmetric around $X = 0$ point. For strong inhomogeneity ($\bar{\lambda} \gg 0$), there is a strong asymmetry of the effective potentials. The effective potentials for negative k_y does not have any global minimum. Therefore, the bound state does not exist for negative k_y . The effective potentials together with the energy eigenvalues for various values of $\bar{\lambda}$ are shown in figure 1. Both the effective potentials get saturated to $(\hbar\omega_c(x_0))^2 \xi_0$ at large X . The zero-energy state ($n = 0$) is always lies inside the potential V_- but outside the potential V_+ . However, all other discrete energy levels ($n > 0$) are lying inside both the potentials $V_{\pm}$, which is expected from the solution of the Dirac-Weyl equation. The number of energy levels decreases as we increases $\bar{\lambda}$. For example,




FIG. 1: (Color online) Plots of the effective potential $V_{\pm}(x)$ (in units of $(\hbar\omega_c)^2$) vs X together with the energy eigenvalues E_n (in units of $\hbar\omega_c$) for $\bar{\lambda} = 0.5$ and $\bar{\lambda} = 0.01$ (only zero-energy level is shown in this case).

there are five energy levels including zero-energy state when $\bar{\lambda} = 0.5$. On the other hand, number of energy levels is quite large when $\bar{\lambda} = 0.01$. There are a finite number of discrete energy levels for a given asymmetric parameter $\bar{\lambda}$. The total number of discrete energy levels including zero-energy level can be calculated easily from the condition that $E_n^2 \leq V_{\pm}(X \rightarrow \infty)$ and it is given by $N = \text{Int}[\xi_0] + 1$. Here, $\text{Int}[b]$ means the integer that is just smaller than b . In figure 1, only the zero-energy level is shown for the $\bar{\lambda} = 0.01$ case.

The probability density distribution of Dirac electrons in a given level n is $\rho_n(x) = \Psi^{(n)\dagger}(x)\Psi^{(n)}(x)$. The probability density distributions ρ_n for different values of n and $\bar{\lambda}$ are shown in figure 2. The velocity operator that follows from the Heisenberg equation is given by $\mathbf{v} = v_F\sigma$. The probability cur-

rent distribution is $J_n(x) = ev_F\Psi^{(n)\dagger}(x)\sigma_y\Psi^{(n)}(x) = -iev_F(\Psi_+^{(n)*}(x)\Psi_-^{(n)}(x) - \Psi_-^{(n)*}(x)\Psi_+^{(n)}(x))$. The probability current distributions for various values of n and $\bar{\lambda}$ are shown in figure 3. The zero-energy state does not carry any current, irrespective of nature of the magnetic field. When $\bar{\lambda} \rightarrow 0$, ρ_n and J_n are symmetric around the point $X = 0$. When $\bar{\lambda} \gg 0$, ρ_n and J_n are strongly asymmetric as is expected from the effective potentials.

In summary, we have obtained exact energy eigenvalues and the corresponding eigenfunctions for a Dirac electron in the presence of an exponentially decaying mag-




FIG. 2: (Color online) Plots of the probability density distribution $\rho_n(x)$ (in units of $1/(L_y l(x_0))$) vs X for various values of n with $\bar{\lambda} = 0.5$ and $\bar{\lambda} = 0.01$.

netic field. We have also provided the probability density and current distributions for each band.

I would like to thank K. Bhattacharya for a discussion. This work was supported by a research grant (Grant No.: IITK/PHY/20080036) of DORD, IIT-Kanpur, India.

-
- [1] K. S. Novoselov, D. Jiang, F. Schedin, T. J. Booth, V. V. Khotkevich, S. M. Morozov, and A. K. Geim, PNAS **102**, 10451 (2005).
 - [2] K. S. Novoselov, A. K. Geim, S. M. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorievs, and A. A. Firsov, Science **306**, 666 (2004).
 - [3] K. S. Novoselov, A. K. Geim, S. M. Morozov, D. Jiang, Y. Zhang, M. I. Katsnelson, I. V. Grigorievs, S. V. Dubonos,

- and A. A. Firsov, Nature **438**, 197 (2005).
- [4] Y. Zhang, Y. -W. Tan, H. L. Stormer, and P. Kim, Nature **438**, 2001 (2005).
- [5] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, arXiv: 0709.1163, to appear in Reviews of Modern Physics.
- [6] C. W. J. Beenakker, arXiv: 0710.3848.
- [7] P. R. Wallace, Phys. Rev. **71**, 622 (1947).




FIG. 3: (Color online) Plots of the probability current density $J_n(x)$ (in units of $ev_F/(L_y l(x_0))$) vs X for various values of n with fixed $\bar{\lambda} = 0.5$ and $\bar{\lambda} = 0.01$.

- [8] M. I. Katsnelson, K. S. Novoselov, and A. K. Geim, *Nature* **2**, 620 (2006); J. M. Pereira Jr., V. Minar, F. M. Peeters, and P. Vasilopoulos, *Phys. Rev. B* **74**, 045424 (2006).
- [9] K. S. Novoselov, Z. Jiang, Y. Zhang, S. V. Morozov, H. L. Stormer, U. Zeitler, J. C. Maan, G. S. Boebinger, P. Kim, and A. K. Geim, *Science* **315**, 1379 (2007).
- [10] A. De Martino, L. Dell'Anna, and R. Egger, *Phys. Rev. Lett.* **98**, 066802 (2007).
- [11] L. Oroszlany, P. Rakya, A. Kormanyos, C. J. Lambert, and J. Cserti, *Phys. Rev. B* **77**, 081403(R) (2008).
- [12] T. K. Ghosh, A. De Martino, W. Hausler, L. Dell'Anna, and R. Egger, *Phys. Rev. B* **77**, 081404(R) (2008).
- [13] W. Hausler, A. De Martino, T. K. Ghosh, and R. Egger, in preparation.
- [14] J. E. Mueller, *Phys. Rev. Lett* **68**, 385 (1992).
- [15] F. M. Peeters and A. Matulis, *Phys. Rev. B* **48**, 15166 (1993).
- [16] I. S. Ibrahim and F. M. Peeters, *Phys. Rev. B* **52**, 17321 (1995).
- [17] B. Y. Gu, W. D. Sheng, X. H. Wang, and J. Wang, *Phys. Rev. B* **56**, 13434 (1997).
- [18] K. Handrich, *Phys. Rev. B* **72**, 161308(R) (2005).
- [19] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics* (Pergamon, Oxford 1965).
- [20] G. Menon and S. Belyi, *Phys. Lett. A* **330**, 33 (2004).